

# MAGNETIC FIELD ENHANCED HYBRIDIZATION OF TARGET MOLECULES TO IMMOBILIZED PROBES

Inventors: Jaime E. Ramírez-Vick of Berkeley and Simon Chin of Saratoga, both of  
California.

## BACKGROUND OF THE INVENTION

### a) Field of the Invention

This invention is related to methods for rapid hybridization of complementary nucleic acid molecules and their use in purification, immunoassays, biosensors, and other biochemical applications.

### b) Description of Related Art

The process by which one member of a pair of nucleic acids (oligonucleotides, DNA, RNA) binds its complementary member is known as hybridization. Since nucleic acid molecules have a very strong preference for their sequence complements, simple mixing of complementary sequences is enough to induce them to hybridize. Hybridization is temperature dependent, so nucleic acid molecules that hybridize strongly at low temperatures can be temporarily separated (denatured) by heating. Hybridization is the basis for the polymerase chain reaction and *in situ* hybridization technologies.

Hybridization is a two step process involving (1) diffusion of the target nucleic acid molecule through the hybridization medium until it reaches an immobilized probe and (2) contact with the probe and binding to its complementary sequence. This process is usually slow because the limited amount of target takes very long to diffuse to the probe surface. The diffusion rate is usually increased by increasing the temperature. Although this increases the hybridization rate, there are limits above which the molecules may be irreparably harmed and the process still requires inordinate amounts of time to complete. Numerous parties have addressed the problem of decreasing cycle time in thermal cyclers and related devices.

15  
20  
25  
30  
35  
40  
45  
50  
55  
60  
65  
70  
75  
80  
85  
90  
95  
100  
105  
110  
115  
120  
125  
130  
135  
140  
145  
150  
155  
160  
165  
170  
175  
180  
185  
190  
195  
200  
205  
210  
215  
220  
225  
230  
235  
240  
245  
250  
255  
260  
265  
270  
275  
280  
285  
290  
295  
300  
305  
310  
315  
320  
325  
330  
335  
340  
345  
350  
355  
360  
365  
370  
375  
380  
385  
390  
395  
400  
405  
410  
415  
420  
425  
430  
435  
440  
445  
450  
455  
460  
465  
470  
475  
480  
485  
490  
495  
500  
505  
510  
515  
520  
525  
530  
535  
540  
545  
550  
555  
560  
565  
570  
575  
580  
585  
590  
595  
600  
605  
610  
615  
620  
625  
630  
635  
640  
645  
650  
655  
660  
665  
670  
675  
680  
685  
690  
695  
700  
705  
710  
715  
720  
725  
730  
735  
740  
745  
750  
755  
760  
765  
770  
775  
780  
785  
790  
795  
800  
805  
810  
815  
820  
825  
830  
835  
840  
845  
850  
855  
860  
865  
870  
875  
880  
885  
890  
895  
900  
905  
910  
915  
920  
925  
930  
935  
940  
945  
950  
955  
960  
965  
970  
975  
980  
985  
990  
995

The actual hybridization reaction represents the most important and central step in the whole process. The hybridization step involves placing the prepared nucleic acid sample in contact with a specific reporter probe, at a set of optimal conditions for hybridization to occur to the target sequence. Hybridization may be performed in any one of a number of formats. For example, multi-sample nucleic acid hybridization analysis has been conducted on a variety of filter and solid support formats (See G. A. Beltz et al., in *Methods in Enzymology*, Vol. 100, Part B, R. Wu, L. Grossman, K. Moldave, Eds., Academic Press, N.Y., Chapter 19, pp. 266-308, 1985). One format, the so-called "dot blot" hybridization, involves the non-covalent attachment of target nucleic acids to filters, which are subsequently hybridized with a radioisotope labeled probe(s). "Dot blot" hybridization gained wide-spread use, and many versions were developed (see M. L. M. Anderson and B. D. Young, in *Nucleic Acid Hybridization--A Practical Approach*, B. D. Hames and S. J. Higgins, Eds., IRL Press, Washington, D.C. Chapter 4, pp. 73-111, 1985). It has been developed for multiple analysis of genomic mutations (D. Nanibhushan and D. Rabin, in EPA 0228075, Jul. 8, 1987) and for the detection of overlapping clones and the construction of genomic maps (G. A. Evans, in U.S. Pat. No. 5,219,726, Jun. 15, 1993).

New techniques are being developed for carrying out multiple sample nucleic acid hybridization analysis on micro-formatted multiplex or matrix devices (e.g., DNA chips) (see M. Barinaga, 253 Science, pp. 1489, 1991; W. Bains, 10 Bio/Technology, pp. 757-758, 1992). These methods usually attach specific DNA sequences to very small specific areas of a solid support, such as micro-wells of a DNA chip. These hybridization formats are micro-scale versions of the conventional "dot blot" and "sandwich" hybridization systems. The micro-formatted hybridization can be used to carry out "sequencing by hybridization" (SBH) (see M. Barinaga, 253 Science, pp. 1489, 1991; W. Bains, 10 Bio/Technology, pp. 757-758, 1992).

U.S. Patent No. 5,849,486 teaches the use of electric fields to accelerate hybridization by bringing the negatively charged target molecules close to the immobilized probes. This greatly accelerates hybridization but requires a restricted flow area due to the tendency of the negatively charged DNA to follow the electric field lines, which do not necessarily cross the complementary target DNA location. In order to overcome this problem, the probe DNA has been immobilized within the pores of a polymer such as agarose. Even though this brings the target DNA and the

probe DNA in close contact, the movement across the restricted pores slows down the probe because of the high hydrodynamic resistance.

This approach is also nonspecific in nature since the movement of the probe DNA in the electric field depends on the molecule's charge, which varies from molecule to molecule depending upon its sequence. Another problem with this approach is that the metal used in the manufacturing of the electrodes, which generate the electric field, may become magnetized.

The object of the present invention is to provide an improved method for the rapid hybridization of nucleic acids on solid surfaces. Several longstanding problems in purification, immunoassays, biosensors, and other biochemical applications are resolved by this invention.

## SUMMARY OF INVENTION

The present invention relates to a method to actively carry out controlled multi-step processing and multiplex reactions. These reactions include, but are not limited to, most molecular biological procedures, such as nucleic acid hybridization, antibody/antigen reaction, and related clinical diagnostics. In addition, the claimed methods are useful for multi-step combinatorial biopolymer synthesis, including, but not limited to, the synthesis of different oligonucleotides or peptides at specific micro-locations on a given device.

The invention provides a method for increasing the hybridization rate of nucleic acids in a nucleic acid assay by labeling nucleic acid molecules with paramagnetic labels and activating a magnetic field inducing rapid migration of the labeled molecules to a solid support, where the labeled molecule hybridizes with its complementary pair. The paramagnetic label may be present on the target nucleic acid or the probe nucleic acid, depending upon whether the target or probe are the immobile phase.

## DETAILED DESCRIPTION OF THE INVENTION

### I. Definitions

Unless indicated otherwise, the terms defined below have the following meanings:

“Biopolymer” refers to biological molecules such as proteins, oligonucleotides, DNA, etc., which are the basis of hybridization, purification, immunoassays, and many other biochemical applications.

5 “Hybridization” refers to binding reaction between complementary partners of biopolymer molecules.

“Ligand” refers to one member of the ligand/receptor binding pair, such as, oligonucleotides, DNA, and proteins.

10 “Nucleic acid molecule” refers to any molecule having at least one nucleic acid moiety, including genomic DNA, cDNA, RNA, PNAs, oligonucleotides, and fragments thereof. The nucleic acid moiety is typically A, C, G, T or U, but may also be another natural or synthetic nucleotide.

“Protein” refers to enzymes, antibodies, and other polypeptides.

15 The invention provides a method for increasing the hybridization rate of nucleic acids in a nucleic acid assay by labeling nucleic acid molecules with paramagnetic labels and activating a magnetic field which induces rapid migration of the labeled molecules to a solid support, where the labeled molecule hybridizes with its complementary pair. The paramagnetic label may be present on the target nucleic acid or the probe nucleic acid, depending upon whether the target or probe are the immobile phase.

20 The solid support is selected from the group consisting of silicon, glass, and metals, preferably soft metals selected from the group consisting of silver, copper, gold, platinum (II), mercury, mercury (II), thallium, cadmium (II), platinum (IV) and palladium (II).

25 The paramagnetic labels may be superparamagnetic particles, having a diameter of from about 1 to about 10 nanometers, or are paramagnetic porphyrins. The paramagnetic labels are attached to the nucleic acid molecules using cleavable conjugating molecules. At least one member of a complementary pair may be labeled with a fluorescent detection molecule.

Paramagnetic agents respond very strongly to magnetic fields, becoming demagnetized after the field is removed. The use of a magnetic field decreases the hybridization time from days to minutes. This prevents the aggregation that would be seen if these paramagnetic agents retained any degree of magnetization.

5 This invention describes a device fabricated using photolithographic techniques that produces localized DC magnetic fields. These fields are localized at multiple locations on a substrate and are surrounded by coils. These locations are microwells connected by channels to distribute the target DNA hybridization solutions.

10 The use of paramagnetic agents to concentrate biomolecules using DC magnetic fields is known in the art. The paramagnetic agents currently used are either polymeric capsules with a paramagnetic iron oxide core, beads in which the paramagnetic material is distributed throughout the polymer material, or particles made-up of the paramagnetic material. These agents range in size from a few micrometers to nanometers. The main problem with the use of these agents for hybridization of DNA is their size, since the paramagnetic agent is thousands of times larger than the DNA molecule. This limits the labeled probe DNA accessibility to the immobilized target DNA on the surface and also saturates the available surface for hybridization.

15 An option to the use of these paramagnetic agents is the use of porphyrin molecules. Molecules of various porphyrins consist of a fundamental skeleton of four rings linked together to form a larger ring. The rings are basically of the pyrrole type, four carbon atoms and one nitrogen atom; depending on the porphyrin, various other atoms are connected to the large ring. Moreover, atoms of various metals positioned in the center of the large ring distinguish different porphyrins. Molecules with Gd(III) as the metal ion are currently used as contrast agents for magnetic resonance imaging. Their widespread use as contrast agents is because of their paramagnetic properties. Their paramagnetic properties and size (equivalent to a fluorescent molecule) makes Gd(III) porphyrins ideal candidates for paramagnetic labels of DNA molecules.

25 The following claims are presented to specifically point out and distinctly claim the invention. All documents mentioned in this disclosure are incorporated herein by reference.